



# Current Report

Oklahoma Cooperative Extension Fact Sheets are also available on our website at:  
<http://osufacts.okstate.edu>

## Foliar Fungicides and Wheat Production in Oklahoma – April, 2012

Dr. Bob Hunger

Extension Wheat Pathologist

Department of Entomology & Plant Pathology

Oklahoma State University

Dr. Jeff Edwards

Extension Wheat Agronomist

Plant & Soil Sciences

Oklahoma State University

**Question: How are the growth stages of wheat described?**


**Answer:** The Feekes' scale is commonly used to describe the growth stages of wheat. This scale describes the growth stages of wheat numerically, and is commonly used to indicate the recommended timing of pesticide applications.

**Question: How much damage can a foliar disease such as leaf rust cause on wheat?**

**Answer:** A foliar disease such as leaf rust causes the most damage when it is severe at heading, flowering or milk, and not as damaging at soft dough or later (Table 1).

**Table 1. Approximate percent loss of yield caused by leaf rust at combinations of leaf rust severity and growth stage of wheat.**

| Growth stage | Severity (%) of leaf rust on the flag leaf |    |    |    |     |
|--------------|--|----|----|----|-----|
|              | 10   | 25 | 40 | 65 | 100 |
| Flowering    | 10   | 15 | 20 | 30 | 35  |
| Milk         | 2  | 5  | 8  | 14 | 20  |
| Soft dough   | 1  | 3  | 4  | 7  | 10  |
| Hard dough   | 1  | 1  | 1  | 3  | 5   |



**Figure 1. The Feekes scale of wheat development.**

Large, E.C. 1954. Growth stages in cereals: Illustration of the Feekes' scale. Plant Pathology 3:128-129.

**Question: When should I apply a fungicide?**

**Answer:** All the fungicides listed in Tables 2 and 3 can be applied up to growth stage 10.5 (heads completely emerged but not yet flowering). In most years, the optimum period for application is from growth stages 9 (flag leaf fully emerged) to 10.5 (heads fully emerged) because application in this range most likely provides protection during the critical times of flowering and milk (Table 1).

**Question: What fungicides are available for use in Oklahoma?**

**Answer:** Many fungicides currently are labeled for use on wheat. A comparison of the relative effectiveness of these fungicides is presented in Table 3. REMEMBER to consult the label for the most current and accurate information.

**Question: What is the potential benefit from using a foliar fungicide?**

**Answer:** More than 20 years of fungicide trials including years with little or no disease and several years with high disease pressure have documented an average yield increase of approximately 10 percent from using fungicides. Such an increase usually justifies fungicide use if the yield potential

and price of wheat are high. Hence, consider the following to assist in deciding whether to apply a fungicide to control a foliar disease (Table 4):

- **Will a foliar fungicide help to regain yield?** The answer to this is "NO!" Foliar fungicides can only help protect the yield potential present at application.
- **What is the yield potential of the wheat?** This should be 30 to 40 bu/acre at a minimum, but can go up or down, depending on the price of wheat.
- **What is the price of wheat?** \$5.00/bu or more is desirable.
- **What is the growth stage of the wheat?** Foliar diseases do the most harm when infection is severe at stages such as heading, flowering and milk. So be sure to apply fungicide before disease is severe.
- **What about a split application of a fungicide?** For example, applying half of the full rate at an early growth stage (for example at GS 6 to 7) and the other half or

**Table 2. Effect of foliar fungicides on grain yield, test weight, and severity of wheat leaf rust and powdery mildew.**

|                            | Growth stage <sup>1</sup> | Yield<br>bu/ac | Test weight<br>lb/bu | Leaf rust |    | Powdery mildew<br>% severity |
|----------------------------|---------------------------|----------------|----------------------|-----------|----|------------------------------|
|                            |                           |                |                      |           |    |                              |
| <b>Stillwater 2011</b>     |                           |                |                      |           |    |                              |
| No treatment               | ----                      | 54             | 57                   | 76        |    | 13                           |
| Headline 208 EC @ 3 oz FB* |                           |                |                      |           |    |                              |
| TwinLine 1.753 EC @ 7 oz   | 6 FB 9                    | 60             | 57                   | 33        | <1 |                              |
| Evito 480 SC @ 2 oz        | 9                         | 59             | 58                   | 46        | 10 |                              |
| Evito 480 SE @ 2 oz        | 10.1                      | 59             | 58                   | 7         | 9  |                              |
| Folicur 3.6 F @ 4 oz       | 10.1                      | 59             | 58                   | 3         | 11 |                              |
| Prosaro 421 SC @ 6.5 oz    | 10.1                      | 64             | 58                   | <1        | 8  |                              |
| Quilt 200 SE @ 10 oz       | 10.1                      | 63             | 58                   | 7         | 5  |                              |
| Quilt 200 SE @ 14 oz       | 10.1                      | 60             | 58                   | 2         | 4  |                              |
| Stratego YLD 418 SC @ 4 oz | 10.1                      | 59             | 58                   | 11        | 7  |                              |
| Tilt 3.6 EC @ 4 oz         | 10.1                      | 60             | 58                   | 33        | 8  |                              |
| HSD (P=0.05)               |                           | 9              | 2                    | 26        |    | 8                            |
| <b>Stillwater 2010</b>     |                           |                |                      |           |    |                              |
| No treatment               | ----                      | 44             | 53                   | 99        |    | 78                           |
| Headline 208 EC @ 3 oz FB  |                           |                |                      |           |    |                              |
| TwinLine 1.753 EC @ 9 oz   | 6 FB 10.1                 | 62             | 55                   | 8         | <1 |                              |
| Evito 480 SC @ 2 oz        | 8                         | 58             | 54                   | 88        | 31 |                              |
| Evito 480 SC @ 2 oz        | 10.1                      | 55             | 54                   | 40        | 8  |                              |
| Quilt® 200 SE @ 10.5 oz    | 10.1                      | 61             | 56                   | 23        | 0  |                              |
| Quilt® 200 SE @ 14 oz      | 10.1                      | 63             | 56                   | 11        | 0  |                              |
| Prosaro 421 SC @ 6.5 oz    | 10.1                      | 61             | 55                   | 18        | <1 |                              |
| Tilt® 3.6 EC @ 4 oz        | 10.1                      | 56             | 55                   | 88        | <1 |                              |
| HSD (P=0.05)               |                           | 9              | 3                    | 21        |    | 24                           |

<sup>1</sup> Growth stage 6=first node visible; 8=flag leaf just emerging; 9=flag leaf fully emerged; 10.1= heads just emerging from boot. FB=followed by.

The North Central Regional Committee on Management of Small Grain Diseases (NCERA-184) has developed the following information on fungicide efficacy for control of certain foliar diseases of wheat for use by the grain production industry in the U.S. Efficacy ratings for each fungicide listed in the table were determined by field testing the materials over multiple years and locations by the members of the committee. Efficacy is based on proper application timing to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a single application of the labeled rate as listed in the table. Table includes most widely marketed products labeled products, and is not intended to be a list of all labeled products.

**Table 3. Efficacy of fungicides for wheat disease control based on appropriate application timing.**

| Fungicide(s)         |                       |                            | Stagonospora       |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|----------------------|-----------------------|----------------------------|--------------------|-------------------|-----------------|-----------------|-----------------|-----------------|-----------------|---------------------------|-----------------|------------------------|
| Class                | Active ingredient     | Product                    | Rate/A<br>(fl. oz) | leaf/<br>mildew   |                 | Septoria        |                 |                 |                 |                           | Head<br>Scab    | Harvest<br>Restriction |
|                      |                       |                            |                    | Powdery<br>mildew | glume<br>blotch | leaf<br>blotch  | Tan<br>spot     | Stripe<br>rust  | Leaf<br>rust    | Stem<br>rust <sup>b</sup> |                 |                        |
| Strobilurin          | Azoxystrobin 22.9%    | Quadris 2.08 SC            | 6.2 - 10.8         | F(G) <sup>a</sup> | VG              | VG              | E               | E <sup>c</sup>  | E               | VG                        | NR              | 45 days                |
|                      | Pyraclostrobin 3.6%   | Headline 2.09 SC           | 6.0 - 9.0          | G                 | VG              | VG              | E               | E <sup>c</sup>  | E               | G                         | NR              | Feekes 10.5            |
|                      | Fluoxastrobin 40.3%   | Evito 480 SC               | 2.0-4.0            | G                 | -- <sup>d</sup> | -- <sup>d</sup> | -- <sup>d</sup> | -- <sup>d</sup> | VG              | -- <sup>d</sup>           | -- <sup>d</sup> | 40 days                |
| Triazole             | Metconazole 8.6%      | Caramba 0.75 SL            | 10.0 - 17.0        | VG                | VG              | -- <sup>d</sup> | VG              | E               | E               | E                         | G               | 30 days                |
|                      | Propiconazole 41.8%   | Tilt 3.6 EC                |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      |                       | PropriMax 3.6 EC           |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      |                       | Bumper 41.8 EC             | 4.0                | VG                | VG              | VG              | VG              | VG              | VG              | P                         | Feekes 10.5     |                        |
|                      | Prothioconazole 41%   | Proline 480 SC             | 5.0 - 5.7          | -- <sup>d</sup>   | VG              | VG              | VG              | -- <sup>d</sup> | VG              | VG                        | G               | 30 days                |
|                      | Tebuconazole 38.7%    | Folicur 3.6 F <sup>e</sup> | 4.0                | G                 | VG              | VG              | VG              | E               | E               | E                         | F               | 30 days                |
|                      | Prothioconazole 19%   |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Tebuconazole 19%      | Prosaro 421 SC             | 6.5 - 8.5          | G                 | VG              | VG              | VG              | E               | E               | E                         | G               | 30 days                |
|                      | Metconazole 7.4%      |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Pyraclostrobin 12%    | TwinLine 1.75 EC           | 7.0 – 9.0          | G                 | VG              | VG              | E               | E               | VG              | NR                        | Feekes 10.5     |                        |
| Mixed mode of action | Propiconazole 11.7%   |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Azoxystrobin 7.0%     | Quilt 200 SC               | 14.0               | VG                | VG              | VG              | E               | E               | VG              | NR                        | Feekes 10.5     |                        |
|                      | Propiconazole 11.7%   |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Azoxystrobin 13.5%    | Quilt Xcel 2.2 SE          | 14.0               | -- <sup>d</sup>   | VG              | -- <sup>d</sup> | -- <sup>d</sup> | -- <sup>d</sup> | VG <sup>f</sup> | -- <sup>d</sup>           | NR              | Feekes 10.5            |
|                      | Propiconazole 11.4%   |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Trifloxystrobin 11.4% | Stratego 250 EC            | 10.0               | G                 | VG              | VG              | VG              | VG              | VG              | NR                        | 35 days         |                        |
|                      | Prothioconazole 10.8% |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Trifloxystrobin 32.3% | Stratego YLD               | 4.0                | VG                | VG              | VG              | VG              | VG              | VG              | NR                        | 30 days         |                        |
|                      | Tebuconazole 22.3%    |                            |                    |                   |                 |                 |                 |                 |                 |                           |                 |                        |
|                      | Trifloxystrobin 22.3% | Absolute 500 SC            | 5.0-7.7            | VG-G              | -- <sup>d</sup> | -- <sup>d</sup> | -- <sup>d</sup> | -- <sup>d</sup> | E               | -- <sup>d</sup>           | NR              | 4 days                 |

<sup>a</sup> Efficacy categories: NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent. Efficacy designation with a second rating in parenthesis indicates greater efficacy at higher application rates.

<sup>b</sup> Efficacy may be significantly reduced if solo strobilurin products are applied after stripe rust infection has occurred.

<sup>c</sup> Insufficient data to make statement about efficacy of this product.

<sup>d</sup> There are several generic products containing tebuconazole but products may not be labeled in all states.

<sup>e</sup> Estimates of fungicide efficacy against stem rust are based on a small number of observations, and may be less reliable than the ratings for other diseases.

This information is provided only as a guide. It is the responsibility of the pesticide applicator by law to read and follow all current label directions. No endorsement is intended for products listed, nor is criticism meant for products not listed. Members or participants in the NCERA-184 committee assume no liability resulting from the use of these products.

a full rate at GS 10 or so. Splitting the application of a fungicide may provide benefit for diseases such as tan spot, septoria/stagonospora and powdery mildew that initiate from fungal inoculum on wheat residue left on the soil surface such as in no-till situations. A split application also may have benefit if a variety is extremely susceptible to these diseases and they are present in the late winter or early spring. However most data indicates that a single application from stages 9 to 10.5 is usually the most beneficial. If a split application is used, the first application should not be made with topdressing as the nitrogen needs to be applied prior to finding nodes at the base of tillers (GS 6 to 7) so the fertilizer moves into the root zone prior to jointing.

- **What diseases are present?** Be sure which foliar fungal diseases are present. Stripe rust can be especially damaging because of its ability to quickly kill entire leaves. Hence, if you are considering a fungicide application to

protect against stripe rust, it is critical to apply the fungicide before the appearance of rust pustules on the flag leaf.

- **What is the disease reaction of the variety?** Refer to the OSU Extension Fact Sheet PSS-2142 "Wheat Variety Comparison Chart," available online at osufacts.okstate.edu or <http://www.wheat.okstate.edu/wheatmanagement/varieties/index.htm>. Some pathogens (e.g., the pathogen that causes wheat leaf rust) can adapt to resistance genes, and a resistant variety may become susceptible when a new race appears.
- **What is the weather forecast?** Hot and dry conditions inhibit further disease development and hasten ripening, while cool and moist conditions promote disease and lengthen the period of time for grain development and filling.

**Table 4. The formulas below can be used to help determine the potential value of a fungicide application. This is a simple cost-benefit evaluation where the yield potential, the price of a bushel of wheat, and the cost of a fungicide can all be easily adjusted.**

| Potential increase                                 | Estimated yield goal | Estimated selling price | Fungicide + app. cost | Potential return on investment |
|--|----------------------|-------------------------|-----------------------|--------------------------------|
| <b>Grain production scenario</b>                   |                      |                         |                       |                                |
| 10%  | X                    | 30 bu/A                 | X                     | \$5.00/bu                      |
| 10%  | X                    | 50 bu/A                 | X                     | \$5.00/bu                      |
| <b>Same scenario for certified seed production</b> |                      |                         |                       |                                |
| 10%  | X                    | 30 bu/A                 | X                     | \$10.00/bu                     |
| 10%  | X                    | 50 bu/A                 | X                     | \$10.00/bu                     |

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0312 GH Revised